

ORIGINAL ARTICLE:

ASSOCIATION BETWEEN DEPRESSION AND SOCIAL NETWORKING SITE USE: A CROSS-SECTIONAL SURVEY AMONG STUDENTS OF A MEDICAL COLLEGE IN PAKISTAN.

Sadaf Riaz¹, Humaira Mahmood², Sahar Riaz³, Nimal Imtiaz⁴.

^{1,2,4}Armed Forces Post Graduate Medical Institute (AFPGMI), National University of Health Sciences (NUMS), Rawalpindi, Pakistan.

³Department of Psychiatry, Beaumont Hospital, North Dublin Mental Health Services (NDMHS), Health Service Executive (HSE), Ireland.

CORRESPONDENCE: DR SADAF RIAZ

Email: sadaf.u2288@gmail.com

Submitted: January 20, 2021

Accepted: October 22, 2023

ABSTRACT

OBJECTIVE

To assess if there is any association between symptoms of depression and social media use among medical students in Pakistan.

STUDY DESIGN

Analytical cross-sectional quantitative research

PLACE AND DURATION OF STUDY

The study was conducted at HITEC Institute of Medical Sciences, Taxilla, Pakistan over a period of about one year, from August 2019 to August 2020.

METHOD

The study participants were medical students. The total sample size was 410. Random sampling was used. Data was collected by using a structured questionnaire. Depressive symptoms were assessed using the MDI (Major Depressive Inventory scale). Time spend on social networking sites was measured using a modified SONTUS scale (Social Network Time Use Scale). The data was analysed with SPSS v21.0 software.

RESULTS

The study found out that there was a significant association between Social Networking Site (SNS) use and depression. A total of 267 (65 %) students reported some form of depression. For SNS use, 163 (40%) were high users and 19 (4.6%) were extremely high users.

CONCLUSION

This research suggests an important relationship between social media usage and depressive symptoms. A high number of students were found to be severely depressed. This reveals a neglected area of mental health and needs to be addressed. Also, almost half of the respondents were high users of social networking sites, while most others were average users. This might indicate that excessive use of SNS might either be a contributing factor or a perpetuating factor for depression in medical students.

KEYWORDS

Depression; Mental Health; Pakistan; Social-Media; Students, Medical.

INTRODUCTION

Increased usage of social networking and the growing mental health issues are raising concerns among professionals all over the world. The incidence of depression is increasing at a gradual pace, affecting the functioning and quality of life of millions of individuals around the globe. The estimated economic burden in the world due to mental disorders in 2010 was about US\$ 8.5 trillion, and this burden is expected to increase to twice this amount by 2030¹. In Pakistan, the rates of mental health disorders are also on the rise, with anxiety and depression ranging from 22-60% of the general population².

Young adults in medical colleges are exposed to various factors during their years of study, and multiple studies have documented high levels of depression, anxiety, and stress among medical students. A study³ examined the stigma among medical students towards depression and suicidal ideation, and found a significant prevalence of depression in medical students; moderate to severe depression in 14.3%, suicidal ideation in third- and fourth- year medical students was seen in 7.9%, and significantly prevailing stigma among medical students regarding depression, anxiety and suicidal ideation. Thus, exploring a cause that might contribute to low mood and depressive symptoms among young medical students is crucial. The future healthcare providers must have optimum health themselves in order to provide care to others, and investigating the issues which might affect the mental health of medical students is necessary to make them into an effective workforce.

Evidence suggests that although a lot of factors contribute to low mood in young adults, the potential influence of social media (SM) use in relation to mental well-being is gaining interest. Zheng et al⁴ looked at the various negative consequences of excessive use of social networking sites on family, personal and professional life, using principles of the cognitive behavioural model and social cognitive theory. A study conducted in New Zealand⁵ established that the prevalence of depression was higher for medical students than the general population, and explored various factors in a medical student's life that might contribute to the higher rates of depression and anxiety, including various assessment tools, selection procedures, students' motivation, characteristics of the student population (such as Type A personality and perfectionism), resilience and the nature of the clinical environment. However, the study did not explore the increased role of the internet, social media and social networking in the daily lives of medical students. This needs to be addressed in order to understand its contribution or relationship to the increasing prevalence of depression among young adults.

Clark et al⁶ discussed how social media sites could have both negative and positive consequences, based on how they were used; social networking sites benefited people when they were used to form meaningful connections and bonds, and harmed people due to social comparison and feelings of isolation. A study⁷ conducted among students from a university in China revealed that the type of social networking sites usage mattered; the study showed that students agreed that 'social' social networking site use was positively related to their wellbeing, while 'entertainment' type social networking site use was not.

In Pakistan, very limited amount of research has been done regarding any connection between Social Networking Site (SNS) use and depressive symptomology. SNS are "mobile- or Internet-based social platforms created and designed to enable users to communicate, collaborate and share content across contacts and communities"⁸. Due to social pressures and easy availability of the internet among the young generation, social media use has increased from functional to problematic usage. This study was conducted keeping in view the lack of available data on SNS use and the presence of depressive symptomatology in medical students in Pakistan, as well as understanding the relationship between them.

Understanding this association will result in better usage of SNS in their capacity to positively affect mental health of future doctors and caregivers. The objective of this study is to look for a relationship between Depression and Social Network use among medical students in Pakistan. This can help improve the mental health of young adults by raising awareness and giving evidence-based recommendations regarding Social Networking Site use.

METHOD

This analytic cross-section study was carried out for a period of 1 year (August 2019 to August 2020) among medical students at HITEC Institute of Medical Sciences in Taxila, Pakistan. Ethical approval was obtained from the ethical review committee of HITEC Institute of Medical Sciences (ERC/21/20, dated 5th June 2020) prior to the study.

The study population was MBBS (medical) students aged 18 to 25 years, from first to fifth years of medical college. A total sample of 410 students who were available on the day of the survey were chosen via random sampling technique with 82 students from each year. Full informed consent was obtained from each participant prior to their involvement in the study. Exclusion criteria included students taking any psychotropic drugs or those who had suffered from a bereavement in the past six months. Those who refused to participate, returned incomplete questionnaires or missed entries were also excluded.

A structured questionnaire was administered, including demographics, the MDI (Major Depressive Inventory) scale⁹, and a modified SONTUS scale (Social Network Time Use Scale). The SONTUS scale was used for assessing the time expended on social networking sites (SNS)¹⁰.

Operational definitions:

SNS, or social networking sites are defined as 'mobile- or internet- based social platforms created and designed to enable users to communicate, collaborate and share content across contacts and communities'. The social networking sites included in this study were Facebook, Instagram, Twitter (X), LinkedIn and Whatsapp.

Depression was defined according to the DSM-5 criteria; 2 or more weeks of either low mood or disinterest/displeasure, along with five or more from the following criteria: depressed mood almost every day, anhedonia almost every day, major weight loss or gain, a deceleration of thought and decrease in body exertion, fatigue or energy loss, feelings of worthlessness or guilt, decreased concentration, persistent thoughts of death. For students who scored high on the MDI, they were provided support and options to link in with a mental health specialist for further care.

Pilot Study

The structured questionnaire was initially validated in a pilot study carried out among 30 respondents in a medical college to check for the feasibility of the study and to check for the reliability and validity of the questionnaire. Content validity is already established as the data collection tool is validated and has been used in various studies. Cronbach's alpha (α)

was used to calculate the internal consistency reliability using SPSS software version 21 and it was found to be 0.816.

Data Collection

After obtaining approval from the institution, students were approached and consent was taken from them. Students were chosen at random, based on their availability on the day of the survey. They were first explained about the context of the survey, the need to assess mood and time spent on social networking sites, and that it would help us in improving care and mental health of medical professionals. The questionnaire was then explained in clear, simple and local language if they were not able to understand properly. They were also allowed to ask questions where ever they were confused or had a query. The questionnaire was filled only after the investigator was sure that the student understood the questions clearly and they understood their right to withdraw from the study at any time.

Statistical Analysis

Analysis of data for this research was done using the Statistical Package of Social Sciences (SPSS) version 21. For descriptive statistics, numerical variables were stated as means and standard deviations, and categorical variables were expressed as frequencies and percentages. Level of significance for rejecting the null hypothesis was 0.05 (less than 0.05 was significant). For inferential statistics, Chi square test was used to look for any significant relationship among independent variables and dependent variables for categorical data.

RESULTS

A total of 410 medical students from the medical college completed the questionnaire. The sample comprised of 34.9% males and 65.1% females, aged between 18- 25 years, with a mean age of 20.95 years (SD 1.716). Approximately 48.3% of the medical students resided in homes and 51.7% in hostels. 20% students were taken from each year of the medical college, i.e., from the first to fifth year.

According to MDI (Major Depressive Inventory Scale), depression was divided into four categories: no depressive symptoms, mild, moderate and severe depressive symptoms. Table 1 shows the number of respondents in each category of depression.

Table 1
Frequency of depression among medical students using MDI scale.

Severity of Depression	Frequency	Percentage
No Depression	143	34.9%
Mild Depression	42	10.2%
Moderate Depression	64	15.6%
Severe Depression	161	39.3%
Total	410	100%

The scale used for SNS use was divided into four categories, ranging from low to extremely high users. See table 2 for results. Almost half the respondents were average users of SNS, followed by almost 40% of the respondents being high users

Table 2
Frequency of Social Network Site Use among medical students using modified SONTUS scale.

SNS use frequency	Frequency	Percentage
Low User	52	12.7%
Average User	176	42.9%
High User	163	39.8%
Extremely High User	19	4.6%
Total	410	100%

A chi-square test of independence was performed to examine the association between depression and social network site use. Almost 30% of the students answered questions 2,3 and 8 (29.0%, 31.7% and 32.9% respectively) i.e., loss of interest in daily activities, lack of energy and strength, and feeling restless or subdued most of the time. About half of the respondents answered all the questions as having all these feelings some of the time (Table 3). The relation between these variables was significant, $\chi^2(9, N = 410) = 41.11, p = <.001$ which tells us that there is a clear association between presence of depression and depression severity and SNS use.

Table 3
Association between Depression and SNS Use.

Severity of Depression	Low user n (%)	Average user n (%)	High user n (%)	Extremely high user n (%)	p-value
No depression	25 (17.5%)	78 (54.5%)	33 (23.1%)	7 (4.9%)	<0.001
Mild depression	5 (11.9%)	22 (52.4%)	15 (35.7%)	0 (0.0%)	
Moderate depression	12 (18.8%)	22 (34.4%)	27 (42.2%)	3 (4.7%)	
Severe depression	10 (6.2%)	54 (33.5%)	88 (54.7%)	9 (5.6%)	

Note: N=frequency and %= percentage.

DISCUSSION

This study reveals a strong correlation between depression and social networking site (SNS) use among young medical undergraduates. While some studies have shown mixed results or no correlation¹¹, these findings align with previous research linking SNS use and depression.

The prevalence of depression in this study was 65.1%, comparable to earlier research. Various studies show that medical students have anxiety, stress, and depression¹². Reported prevalence rates include 51.3% among Indian medical students, 83.4% among Saudi students¹³, and 40.9% in Pakistan¹⁴. A separate study in Pakistan¹⁵ found depressive symptoms in 45.5% of 437 medical students, similar to this study's outcome. Another 2017

study from a private Pakistani medical college, reported depression and anxiety symptoms in 51.46% of students¹⁶, while a more recent study in Lahore¹⁷ found a 75% prevalence among 533 students.

However, a meta-analysis of 195 studies across 43 countries before 2016 reported a much lower depression prevalence of 27.2%¹⁸. The past decade has seen a tenfold increase in social media use, with nearly 90% of young adults (18–29) using it by 2015. A meta-analysis suggests approximately 75% of medical students use SNS¹⁹, and Pakistani students are no exception.

In this study, all participants reported SNS use. Among the 410 respondents, 12.7% were low users, 42.9% were average users, 39.8% were high users, and 4.6% extremely high users. Nearly half were average users, followed by high users. A study in East India found SNS usage among medical undergraduates to be 88.6%²⁰, supporting this study's findings.

The impact of SNS use is multifaceted. Previous research links it to depression, anxiety, stress, mood swings, eye pain²¹, and internet addiction²². Consistent with this study, research indicates that higher SNS use correlates with increased depression risk, largely due to excessive time spent online. Frequent social media use has been associated with mental health issues, including "Facebook Depression Phenomenon," which stems from prolonged online engagement, social comparison, and low mood. Studies suggest SNS exposure can negatively affect self-esteem and lead to adverse psychological outcomes²³. Other effects include reduced in-person social interactions²⁴, decreased attention span, and increased cyberbullying risks, all contributing to poor mental well-being.

Conversely, some studies report no association between SNS use and depression²⁵. These mixed findings highlight the complexity of measuring this relationship, as SNS use is multifaceted. Regardless of directionality, these results are important for public health professionals and clinicians. Further research is needed to explore this relationship in greater depth and draw evidence-based conclusions.

It is also important to recognize that SNS interactions vary widely. This study measured total time spent on SNS rather than specific interactions. Additionally, prior research has focused mostly on Facebook, while this study examined SNS use more broadly in relation to depression.

CONCLUSION

This research explored, in particular, the impact of social network site use on mental health and found important evidence to suggest a positive correlation between the use of social networking sites and depression. A high number of medical students were also found to have depressive symptoms in this study. This reveals a neglected area of mental health of the future health professionals and needs to be addressed. These results provide us with insight and fundamental data about mental health to establish strategies to reduce the burden among medical students.

Further studies with a larger sample size from various other medical colleges are warranted to clarify and contribute to the results of this present research. Also, in order to find out the

cause and effect, a longitudinal study needs to be conducted to study causality rather than knowing simple association. Apart from SNS use, more factors need to be explored which contribute to depression among medical students. A lot of studies have already identified a few, with stress of studies and exams, increased workload etc. Lastly, as far as the current post COVID-19 pandemic world is concerned, measures like lockdowns and social distancing increased SNS use globally, and the effects of that need to be studied on students.

CONFLICT OF INTEREST

The authors declare no conflict of interest

FUNDING

The authors received no funding for this research

DISCLOSURE

This article has been written as a part of the thesis for 'Masters in Public Health' at AFGMI.

REFERENCES

1. Trautmann S, Rehm J, Wittchen HU. The economic costs of mental disorders: Do our societies react appropriately to the burden of mental disorders?. *EMBO reports*. 2016 Sep;17(9):1245-9.
2. Ahmed B, Enam SF, Iqbal Z, Murtaza G, Bashir S. Depression and anxiety: a snapshot of the situation in Pakistan. *International Journal of Neuroscience and Behavioral Science*. 2016;4(2):32.
3. Schwenk TL, Davis L, Wimsatt LA. Depression, stigma, and suicidal ideation in medical students. *Jama*. 2010 Sep 15;304(11):1181-90
4. Zheng X, Lee MK. Excessive use of mobile social networking sites: Negative consequences on individuals. *Computers in Human Behavior*. 2016 Dec 1;65:65-76.
5. Moir F, Yelder J, Sanson J, Chen Y. Depression in medical students: current insights. *Advances in medical education and practice*. 2018 May 7:323-33.
6. Clark JL, Algoe SB, Green MC. Social network sites and well-being: The role of social connection. *Current Directions in Psychological Science*. 2018 Feb;27(1):32-7.
7. Wang JL, Jackson LA, Gaskin J, Wang HZ. The effects of social networking site (SNS) use on college students' friendship and well-being. *Computers in Human Behavior*. 2014 Aug 1;37:229-36.
8. Thurairaj S, Hoon EP, Roy SS, Fong PW. Reflections of Students language Usage in Social Networking Sites: Making or Marring Academic English. *Electronic Journal of E-learning*. 2015 Apr 1;13(4):pp301-315.
9. Bech P, Timmerby N, Martiny K, Lunde M, Søndergaard S. Psychometric evaluation of the Major Depression Inventory (MDI) as depression severity scale using the LEAD (Longitudinal Expert Assessment of All Data) as index of validity. *BMC psychiatry*. 2015 Dec;15(1):1-7.

10. Olufadi Y. Social networking time use scale (SONTUS): A new instrument for measuring the time spent on the social networking sites. *Telematics and Informatics*. 2016 May 1;33(2):452-71.
11. Moreno MA, Jelenchick L, Koff R, Eickhoff J. Depression and internet use among older adolescents: An experience sampling approach. *Psychology*. 2012 Sep 25;3(09):743.
12. Waqas A, Khan S, Sharif W, Khalid U, Ali A. Association of academic stress with sleeping difficulties in medical students of a Pakistani medical school: a cross sectional survey. *PeerJ*. 2015 Mar 12;3:e840.
13. Alharbi H, Almalki A, Alabdan F, Haddad B. Depression among medical students in Saudi medical colleges: a cross-sectional study. *Advances in medical education and practice*. 2018 Dec 4:887-91.
14. Rizvi F, Qureshi A, Rajput AM, Afzal M. Prevalence of depression, anxiety and stress (by DASS scoring system) among medical students in Islamabad, Pakistan. *Br J Med Med Res*. 2015;8(1):69-75
15. Hashmi AM, Aftab MA, Naqvi SH, Sajjad W, Mohsin M, Khawaja IS. Anxiety and depression in Pakistani medical students: a multicenter study. *Health Med*. 2014;8(7):813-20.
16. Azad N, Shahid A, Abbas N, Shaheen A, Munir N. Anxiety and depression in medical students of a private medical college. *Journal of Ayub Medical College Abbottabad*. 2017 Jan 25;29(1):123-7.
17. Zafar U, Daud S, Khalid A. Determinants of depression among undergraduate medical students of a private medical college in Lahore. *JPMMA*. 2020 Jan 14;34:2019.
18. Rotenstein, L. S., Ramos, M. A., Torre, M., Segal, J. B., Peluso, M. J., Guille, C., ... & Mata, D. A. (2016). Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *Jama*, 316(21), 2214-2236.
19. Guraya SY. The usage of social networking sites by medical students for educational purposes: a meta-analysis and systematic review. *North American journal of medical sciences*. 2016 Jul;8(7):268.
20. Lahiry S, Choudhury S, Chatterjee S, Hazra A. Impact of social media on academic performance and interpersonal relation: A cross-sectional study among students at a tertiary medical center in East India. *Journal of education and health promotion*. 2019;8.
21. Madaiah M, Seshaiyengar CT, Suresh P, Munipapanna S, Sonnappa SD. Study to assess the effects of social networking sites on medical college students. *International J Community Med Public Health*. 2017;3(5):1204-8.
22. Ahmer Z, Tanzil S. Internet addiction among social networking sites users: Emerging mental health concern among medical undergraduates of Karachi. *Pakistan Journal of medical sciences*. 2018 Nov;34(6):1473.
23. Barman L, Mukhopadhyay DK, Bandyopadhyay GK. Use of social networking site and mental disorders among medical students in Kolkata, West Bengal. *Indian journal of psychiatry*. 2018 Jul;60(3):340
24. Yoon S, Kleinman M, Mertz J, Brannick M. Is social network site usage related to depression? A meta-analysis of Facebook–depression relations. *Journal of affective disorders*. 2019 Apr 1;248:65-72.

25. Alfasi Y. The grass is always greener on my Friends' profiles: The effect of Facebook social comparison on state self-esteem and depression. Personality and Individual Differences. 2019 Sep 1;147:111-7.

AUTHOR(S) CONTRIBUTION/UNDERTAKING FORM

Sr. #	Author(s) Name	Author(s) Affiliation	Contribution
1	Dr Sadaf Riaz	AFPGMI (Armed Forces Post Graduate Medical Institute), NUMS (National University of Health Sciences), Rawalpindi	Conception of research, data acquisition, analysis and interpretation of data, critical evaluation of article
2	Dr Humaira Mahmood	AFPGMI (Armed Forces Post Graduate Medical Institute), NUMS (National University of Health Sciences), Rawalpindi	Supervision of research, assisting in data analysis, providing critical feedback
3	Dr Sahar Riaz	Department of Psychiatry, Beaumont Hospital, North Dublin Mental Health Services (NDMHS), Health Service Executive (HSE), Ireland	Conception and design of research, drafting the article, critical evaluation of research
4	Dr Nimbal Imtiaz	AFPGMI (Armed Forces Post Graduate Medical Institute), NUMS (National University of Health Sciences), Rawalpindi	Assisting in data acquisition, critical evaluation of article, assisting in data analysis